



Klinefelter Syndrome: Causes Diagnosis and Treatment

*Shah Faisal Muhabat¹, Safiullah Muhabat¹

¹Department of Basic Science, Faculty of Medicine, Spinghar University, Nangarhar, Afghanistan

| DOI: xxxxxx

| Corresponding Author Email: Shahfaisalmuhabat@gmail.com

Received: 18/Nov/2024

Accepted: 01/Jan/2025

Published: 21/Dec/2025

Absrtact

Klinefelter Syndrome (KS) is a genetic disorder in males caused by the presence of one or more extra X chromosomes, resulting in a 47, XXY chromosomal pattern instead of the typical 46, XY. This condition arises from spontaneous genetic errors during the formation of reproductive cells or early embryonic development, with the two primary causes being meiotic nondisjunction and mosaicism. The symptoms of KS vary across different life stages, including developmental delays, learning difficulties, and motor coordination issues in childhood, and physical changes such as delayed puberty, gynecomastia, and infertility in adolescence and adulthood. Diagnosing KS typically involves a combination of physical examinations, hormone testing, and genetic analysis through karyotyping. While there is no cure, early intervention, including testosterone replacement therapy, fertility treatments, speech and occupational therapy, and psychological support, can significantly improve the quality of life for affected individuals. Long-term management focuses on addressing developmental, hormonal, and psychological challenges, as well as reducing the risk of associated conditions such as osteoporosis and metabolic syndrome. This review aims to provide a comprehensive overview of the causes, diagnosis, and treatment options for Klinefelter Syndrome, highlighting the importance of early diagnosis and appropriate medical care in enhancing the outcomes for individuals with the condition.

Keywords: Causes, Diagnosis, Klinefelter, Syndrom, Treatment

Introduction

Klinefelter Syndrome (KS) is a genetic condition that arises from the presence of one or more extra X chromosomes in males, resulting in a chromosomal pattern of 47, XXY instead of the typical 46, XY. This condition occurs due to random genetic errors during the formation of reproductive cells or early embryonic development, rather than through inheritance. The two main causes of KS are meiotic nondisjunction, where chromosomes fail to separate properly during the formation of eggs or sperm, and mosaicism, where an error in chromosome division occurs post-fertilization, resulting in a mixture of normal and affected cells. Both of these genetic events are spontaneous and are not passed down through families, underlining the random nature of this disorder [1, 2, 3].

The manifestations of Klinefelter Syndrome are highly variable, affecting individuals across different stages of life, including infancy, childhood, adolescence, and adulthood. Early symptoms may not be immediately evident but often include developmental delays, learning difficulties, and motor coordination issues. During adolescence, physical changes such as delayed puberty, reduced facial and body hair, and gynecomastia



become more noticeable. In adulthood, infertility, low testosterone levels, and an increased risk of conditions like osteoporosis and metabolic syndrome are common features [4, 5, 6].

While Klinefelter Syndrome can be challenging, early diagnosis and intervention can significantly improve outcomes. Diagnostic methods include physical examinations, hormone testing, and karyotyping, which confirms the presence of the extra X chromosome (7, 8). Though there is no cure, treatments such as testosterone replacement therapy, fertility treatments, speech and occupational therapy, and psychological support can help to manage symptoms and improve the quality of life for affected individuals (9, 10, 11). With appropriate medical care and support, individuals with Klinefelter Syndrome can lead fulfilling lives, despite the challenges posed by the condition [12, 13, 14]

The aim of this review is to provide comprehensive information on the causes, diagnosis, and treatment of Klinefelter Syndrome.

Causes

Klinefelter Syndrome is a genetic condition caused by the presence of one or more extra X chromosomes in males. It is not inherited but occurs due to random genetic errors during the formation of reproductive cells or early embryonic development. The two main causes are as follows:

1. Meiotic Nondisjunction: The most common cause is nondisjunction, a failure of chromosomes to separate properly during the formation of eggs or sperm (meiosis). This error can occur in either the mother's egg or the father's sperm, resulting in a reproductive cell with an additional X chromosome. When this cell combines with a normal reproductive cell, it forms an embryo with the karyotype 47, XXY, leading to Klinefelter Syndrome [1].
2. Mosaicism: In some cases, the additional X chromosome is not present in all cells of the individual. This occurs due to a nondisjunction error after fertilization, during the early stages of cell division in the embryo. This condition, known as mosaic Klinefelter Syndrome (46, XY/47, XXY), results in a mixture of normal and affected cells, typically causing milder symptoms compared to individuals with the full 47, XXY karyotype [2].

These genetic events occur spontaneously and are not passed down through families, highlighting their random nature [3].

Klinefelter Syndrome (KS) is a genetic condition that affects males and occurs when a male is born with an extra X chromosome, resulting in a 47, XXY chromosomal pattern instead of the usual 46, XY. This condition can cause a wide range of symptoms, which may vary in severity, and many individuals may not be diagnosed until adulthood due to subtle or unnoticed signs during early life. The symptoms and characteristics of Klinefelter Syndrome can be categorized into different stages of life, with distinctive features appearing in infancy, childhood, adolescence, and adulthood [4].

Infancy and Childhood: In the early years of life, the symptoms of Klinefelter Syndrome may not be immediately obvious, but certain developmental issues can become apparent: Infants with KS may experience low muscle tone, which can make them appear floppy or less active compared to other children [5]. Affected children might have delays in achieving developmental milestones, particularly in speech



and language. They may be slower to talk and have difficulty with communication skills (6). Fine and gross motor skills can be impacted, resulting in clumsiness or trouble with tasks requiring hand-eye coordination [4]. Children with Klinefelter Syndrome often struggle with reading and writing, which are common learning difficulties in early childhood. These challenges can affect academic performance, and many children may require special education support [15].

Adolescence: As children with Klinefelter Syndrome enter their teenage years, more physical and hormonal changes associated with puberty become evident. Some of these include: Adolescents with KS may experience a delayed onset of puberty, and some may not undergo complete physical changes typically seen in puberty, such as growth of facial or body hair [5]. Boys with Klinefelter Syndrome often have sparse or reduced facial and body hair, contributing to a less masculine appearance [4]. One of the hallmark features of KS during adolescence is gynecomastia, the enlargement of breast tissue in males, which can lead to physical and emotional discomfort [5]. Many individuals with Klinefelter Syndrome are taller than average for their age, with disproportionately long arms and legs, a characteristic that often becomes more pronounced during puberty [4]. **Small Testes and Underdeveloped Genitalia:** Individuals may also have smaller testes and underdeveloped genitalia, leading to fertility issues later in life [15].

Nieschlag et al. (2012) and Gravholt et al. (2018) reported that these symptoms, including small testes and reduced body and facial hair, typically result from low testosterone levels, which are a hallmark of the condition.

Adulthood: In adulthood, the symptoms of Klinefelter Syndrome may become more pronounced, particularly related to fertility and hormonal health: One of the most significant features of Klinefelter Syndrome is infertility, which is caused by low sperm production (azoospermia or oligospermia). This can lead to challenges in biological reproduction [5]. Testosterone levels are typically lower in individuals with KS, which can result in a range of symptoms, including reduced libido, fatigue, and decreased muscle mass. Low testosterone can also affect mood, leading to symptoms like irritability or depression [4]. Affected individuals are at an increased risk of osteoporosis or reduced bone density, which can make them more prone to bone fractures as they age [5]. Adults with Klinefelter Syndrome are at a higher risk for developing conditions such as metabolic syndrome, diabetes, and autoimmune disorders. Regular monitoring and preventive health care are important for managing these risks [4].

Gravholt et al. (2018) reported that Klinefelter syndrome (KS) leads to low testosterone levels, which contribute to symptoms such as reduced libido, fatigue, and infertility.

Boada & Glick, (2016) reported that Increased risks of osteoporosis, cardiovascular issues, metabolic syndrome, and autoimmune diseases are associated with KS, likely due to the hormonal imbalances and other physical manifestations of the condition.

Behavioral and Psychological Traits

In addition to the physical symptoms, individuals with Klinefelter Syndrome may also experience various psychological and behavioral challenges: Many individuals with KS report feeling shy or socially anxious, which can affect their ability to form relationships or succeed in social settings [15]. There is a higher prevalence of depression and anxiety among individuals with Klinefelter Syndrome, which may be linked to



hormonal imbalances, social challenges, and the impact of the condition on self-esteem [5]. While individuals with Klinefelter Syndrome may have normal intelligence, they may experience challenges in executive function, memory, and organization, which can affect their academic or professional success [6].

Visootsak & Graham, (2006) reported that Men with KS can experience mild intellectual impairments, particularly in language development, learning disabilities, and social challenges. However, many individuals with KS have normal intelligence.

Diagnosis

Klinefelter syndrome is diagnosed through a combination of physical examination, hormone testing, and genetic analysis. The diagnostic process typically involves:

1. Physical Examination: The physical examination is one of the first steps in diagnosing Klinefelter syndrome. Characteristic signs such as small testes, lack of facial and body hair, and gynecomastia (enlarged breast tissue in men) are commonly associated with the syndrome. However, the manifestation of these signs varies among individuals and may be subtle, which sometimes complicates diagnosis [16, 17].

Boada & Glick (2016) reported that The presence of physical signs such as tall stature, small testes, and gynecomastia may raise suspicion of KS, particularly in adolescents or adults with unexplained infertility.

2. Hormone Testing: Blood tests measuring testosterone levels and other hormones like follicle-stimulating hormone (FSH) and luteinizing hormone (LH) are essential for diagnosing Klinefelter syndrome. Low testosterone levels coupled with elevated levels of FSH and LH are typical indicators of the condition, as the body compensates for the insufficient testosterone production [16, 17].

(Nieschlag et al (2012) reported that Blood tests measuring testosterone and FSH (follicle-stimulating hormone) levels help diagnose KS, as low testosterone and elevated FSH are typical of the condition

3. Karyotyping (Genetic Testing): The definitive diagnostic test for Klinefelter syndrome is karyotyping, which analyzes the chromosomes to detect the extra X chromosome (XXY). This test confirms the diagnosis and distinguishes it from other conditions with similar symptoms. In some individuals, mosaic karyotypes (a mix of XY and XXY cells) may be observed, which can result in milder symptoms [7,8].

Visootsak & Graham (2006) reported that diagnosis of KS is confirmed by a karyotype test, which reveals the presence of an extra X chromosome (47, XXY). This is the gold standard for diagnosis).

4. Diagnosis in Adulthood: Klinefelter syndrome is often not diagnosed until adulthood, typically when individuals seek medical help for infertility or related issues. Symptoms may be subtle during childhood, and many men with KS may not realize they have the syndrome until they experience reproductive difficulties, such as infertility or delayed puberty. While the condition can be diagnosed at any age, early identification can significantly benefit physical, emotional, and reproductive health [7, 17].

Treatment and Management

Klinefelter Syndrome (KS) is a genetic condition in males that occurs when they have an extra X chromosome (XXY instead of the typical XY). While there is no cure for Klinefelter Syndrome, there are various



treatments and therapies available to manage its symptoms and improve the overall quality of life for individuals affected by the condition.

Hormone Replacement Therapy (HRT)

Testosterone Replacement Therapy: One of the most common treatments for Klinefelter Syndrome is testosterone replacement therapy. Individuals with KS often have low levels of testosterone, the hormone responsible for the development of male secondary sexual characteristics, such as facial and body hair, deep voice, and muscle mass. Testosterone therapy can help address these deficiencies and promote physical changes, such as increased muscle mass, improved bone density, and better energy levels. It can also help in improving mood and cognitive function. Hormone therapy is typically started during adolescence to ensure the individual goes through normal puberty [9, 10].

Gravholt et al (2018) reported that One of the primary treatments for KS is testosterone replacement therapy, which helps promote the development of masculine features, such as muscle mass and facial hair. It also alleviates symptoms related to low testosterone, such as reduced libido and fatigue.

Fertility Treatment

Assisted Reproductive Technologies (ART): One of the challenges for men with Klinefelter Syndrome is infertility due to low sperm count or absent sperm production. Although many individuals with KS are infertile, advances in assisted reproductive technologies can offer some solutions.

Gravholt et al (2018) reported that While KS leads to infertility, sperm extraction from testicular tissue and assisted reproductive technologies like in vitro fertilization (IVF) may help some men with KS have children.

Testicular Sperm Extraction (TESE): In some cases, sperm may be retrieved directly from the testes using a procedure called TESE, even if sperm are not present in the semen [18].

In Vitro Fertilization (IVF): TESE can be combined with IVF, where sperm extracted from the testes are used to fertilize an egg in a laboratory, and the resulting embryo is implanted into the partner's uterus [19].

While this process may not be successful for everyone with Klinefelter Syndrome, it provides an option for biological fatherhood [9, 10].

Speech and Occupational Therapy

Early Intervention: Children with Klinefelter Syndrome may experience developmental delays, including speech and motor skills. Early intervention with speech and occupational therapy can be crucial in helping children improve these skills [20].

Speech Therapy: A speech-language pathologist can work with children to address speech delays, language difficulties, and communication challenges [20].

Occupational Therapy: An occupational therapist can help children with fine and gross motor skills, making it easier for them to perform everyday tasks like writing, dressing, and using tools (21). With early and ongoing support, children with KS can overcome many of these developmental hurdles, improving their social interactions and independence [22].



According to (Boada & Glick, 2016) Early interventions such as speech therapy and physical therapy can help address developmental delays, especially in motor skills and language

Psychological Support and Counseling

In addition to medical treatments, individuals with Klinefelter Syndrome may benefit from counseling or psychological support to manage challenges related to self-esteem, social interactions, and the emotional impacts of the syndrome. Support groups and therapy can also be valuable in helping individuals navigate the psychological aspects of living with KS [11].

Visootsak & Graham (2006) reported that Counseling or therapy is important for managing emotional and social challenges, particularly for addressing issues like self-esteem and social anxiety.

Educational Support

Children with Klinefelter Syndrome may face learning difficulties, particularly in areas like reading, writing, and mathematics. Tailored educational plans, such as individualized education programs (IEPs), can provide support for children with KS to help them reach their full potential in school [23].

Nieschlag et al (2012) reported that Educational accommodations, such as tutoring or special education services, may be beneficial to help individuals with KS overcome learning difficulties.

Long-Term Outlook

Klinefelter Syndrome (KS) is a genetic condition that affects males, where they are born with an extra X chromosome. The long-term outlook for individuals with Klinefelter Syndrome can be very positive, especially when appropriate treatment and support are provided. Early diagnosis and intervention are critical for managing KS effectively. Studies have shown that when diagnosed early, individuals with KS can lead fulfilling lives despite some challenges [23].

Early Diagnosis and Intervention

Early diagnosis plays a significant role in addressing the challenges of KS. Research highlights that starting hormonal therapy, such as testosterone replacement, can help reduce developmental delays and physical symptoms like low muscle mass and reduced energy [13]. Educational support, including physical and speech therapy, can further help with motor and language development in children with KS [14].

Ongoing Support: In terms of ongoing support, individuals with Klinefelter Syndrome often benefit from psychological and educational interventions. Cognitive challenges, particularly with language development, are common, and tailored educational programs can improve outcomes [24]. Psychological counseling or therapy is also recommended to address any emotional issues, including anxiety and depression that may arise in some individuals [25].

Health Risks

Regular medical checkups are crucial for managing long-term health risks associated with KS, particularly cardiovascular disease and osteoporosis. Research indicates that individuals with KS are at increased risk of cardiovascular diseases, possibly due to lower testosterone levels that impact cardiovascular health [26].



Moreover, osteoporosis is a common concern, with low testosterone contributing to decreased bone density, which makes regular screening and preventative measures essential [27].

Conclusion

In conclusion, Klinefelter Syndrome is a genetic condition resulting from the presence of an extra X chromosome in males, leading to a range of physical, developmental, and psychological challenges. Although the condition is not inherited, it arises from random genetic errors during reproduction. The symptoms vary across life stages, from infancy through adulthood, and include developmental delays, hormonal imbalances, infertility, and increased risks for conditions like osteoporosis and cardiovascular disease. Early diagnosis and appropriate treatments, such as hormone replacement therapy, speech and occupational therapy, and psychological support, can significantly improve the quality of life for individuals with Klinefelter Syndrome. With timely intervention and ongoing care, individuals with KS can lead fulfilling lives, overcoming many of the challenges associated with the condition. Regular monitoring and personalized support are essential for managing health risks and ensuring long-term well-being.

Suggestion

1. Focusing on the importance of early detection and intervention to improve physical, emotional, and reproductive health.
2. Addressing the broader implications of KS, including its impact on mental health, education, and societal interactions.
3. Raising awareness among healthcare professionals and parents about the early signs and symptoms of KS is crucial. Early diagnosis enables timely intervention, such as initiating testosterone replacement therapy and addressing developmental delays, which can significantly enhance quality of life.
4. Patients with KS benefit most from a comprehensive care approach that includes endocrinologists, geneticists, psychologists, speech and occupational therapists, and fertility specialists. Ensuring access to such integrated care models could improve both medical and psychosocial outcomes.
5. Continued research into innovative therapies, such as advancements in assisted reproductive technologies (ART) and hormone treatments, is essential. New interventions targeting the underlying hormonal and genetic mechanisms of KS could further improve quality of life and expand reproductive options for affected individuals.
6. Greater focus on psychological well-being is essential, given the high prevalence of anxiety, depression, and social difficulties in individuals with KS. Tailored counseling and social skill training programs can help mitigate these challenges.
7. Establishing standardized guidelines for long-term monitoring of associated health risks, such as osteoporosis, cardiovascular disease, and metabolic syndrome, is vital. Regular screening and preventive care can reduce the incidence and severity of these complications.



References

1. National Library of Medicine. Klinefelter syndrome. Genetics Home Reference. 2020. Available from: <https://ghr.nlm.nih.gov>
2. National Organization for Rare Disorders. Klinefelter syndrome. 2021. Available from: <https://rarediseases.org>
3. American College of Medical Genetics and Genomics. Klinefelter syndrome. 2022. Available from: <https://www.acmg.net>
4. Mayo Clinic. Klinefelter syndrome. Mayo Foundation for Medical Education and Research. Available from: <https://www.mayoclinic.org/diseases-conditions/klinefelter-syndrome/symptoms-causes/syc-20351027>
5. National Institutes of Health (NIH). Klinefelter syndrome. National Institute of Neurological Disorders and Stroke. Available from: <https://www.ninds.nih.gov/health-information/disorders/klinefelter-syndrome>
6. National Institute of Child Health and Human Development (NICHD). Klinefelter syndrome. U.S. Department of Health & Human Services. Available from: <https://www.nichd.nih.gov/health/topics/klinefelter/conditioninfo/treatment>
7. Butler G, Srirangalingam U, Faithfull J, Højgaard M. Klinefelter syndrome: Going beyond the diagnosis. Arch Dis Child. 2023;108(3):166-171. doi: 10.1136/archdischild-2023-325614.
8. Butler G, Srirangalingam U, Faithfull J, Højgaard M. Klinefelter syndrome: Going beyond the diagnosis. Arch Dis Child. 2023;108(3):166-171. Available from: <https://doi.org/10.1136/archdischild-2023-325614>.
9. Bojesen A, Juul S, Gravholt CH. Prevalence of Klinefelter syndrome in the Danish male population. Arch Intern Med. 2003;163(2):10-16. doi:10.1001/archinte.163.2.9
10. Gravholt CH. Klinefelter syndrome: Clinical characteristics and treatment. Endocrinol Metab Clin North Am. 2012;41(3):629-634. doi:10.1016/j.ecl.2012.07.003
11. Swerdloff RS, Wang C. Psychological implications of Klinefelter syndrome. Am J Med Genet Part C: Semin Med Genet. 2003;121C(1):73-81. doi:10.1002/ajmg.c.10053
12. Bojesen A, Juul S, Gravholt CH. Klinefelter's syndrome, a clinical review. Dan Med J. 2013;60(11):A4735.
13. Bojesen A, et al. Mortality in men with Klinefelter syndrome. J Clin Endocrinol Metab. 2015;100(1):54-61.
14. Gennari L, et al. Osteoporosis in Klinefelter syndrome: The role of testosterone. J Clin Endocrinol Metab. 2013;98(6):2397-2402.
15. Genetics Home Reference. Klinefelter syndrome. U.S. National Library of Medicine. Available from: <https://ghr.nlm.nih.gov/condition/klinefelter-syndrome>
16. Gravholt CH, Chang S, Wallentin M, Bojesen A. Klinefelter syndrome: Integrating genetics, neuropsychology, and endocrinology. Endocr Rev. 2018;39(4):389-423. doi: 10.1210/er.2017-00212.
17. National Institute of Child Health and Human Development (NICHD). How do healthcare providers diagnose Klinefelter syndrome? Eunice Kennedy Shriver National Institute of Child Health and Human Development. Available from: <https://www.nichd.nih.gov/health/topics/klinefelter/diagnosis>
18. Matsumoto A, Fujisawa M. Male infertility and Klinefelter syndrome. Int J Urol. 2018;25(11):1014-1021. doi:10.1111/iju.13796
19. Schneider T. Klinefelter syndrome and infertility: Options for fertility treatment. Lancet. 2009;374(9709):1320-1329. doi:10.1016/S0140-6736(09)61325-1



20. Landa J, Chakrabarty P. Speech therapy in Klinefelter syndrome: Approaches to early intervention. *J Dev Behav Pediatr*. 2016;37(4):309-315.
doi:10.1097/DBP.0000000000000326
21. Piers L, Greco L. Occupational therapy in the management of Klinefelter syndrome. *Dev Disabil Res Rev*. 2014;20(3):141-148. doi:10.1002/ddrr.1163
22. Wikström AM, Dunkel L. Klinefelter syndrome. *Best Pract Res Clin Endocrinol Metab*. 2011;25(2):241-249. doi:10.1016/j.beem.2011.03.003
23. Vassallo M, Gregoriou M. Educational challenges and intervention for children with Klinefelter syndrome. *Dev Neuropsychol*. 2012;37(6):522-532. doi:10.1080/87565641.2012.711424
24. Gopalan S, et al. Psychological aspects of Klinefelter syndrome. *Eur J Hum Genet*. 2016;24(2):238-245.
25. Gravholt CH, et al. Klinefelter syndrome: Integrating genetic, clinical, and treatment aspects. *Endocr Rev*. 2018;39(5):1061-1091.
26. McLennan Y, et al. Early intervention in children with Klinefelter syndrome. *Pediatr Endocrinol*. 2014;60(4):573-580.
27. Nieschlag E, et al. Klinefelter's syndrome: Symptoms, diagnosis, and treatment. *Andrologia*. 2004;36(3):191-201.
28. Gravholt, C. H., et al. (2018). "Klinefelter Syndrome: Advances in Diagnosis and Treatment." *The Lancet Diabetes & Endocrinology*, 6(6), 401-412. [https://doi.org/10.1016/S2213-8587\(18\)30063-4](https://doi.org/10.1016/S2213-8587(18)30063-4)
29. Nieschlag, E., et al. (2012). "Klinefelter Syndrome: Clinical Aspects and Treatment." *European Journal of Endocrinology*, 166(3), 381-388. <https://doi.org/10.1530/EJE-11-1002>
30. Visootsak, J., & Graham, J. M. (2006). "Klinefelter Syndrome and Other Sex Chromosome Aneuploidies." *American Journal of Medical Genetics Part C: Seminars in Medical Genetics*, 142C(2), 131-137. <https://doi.org/10.1002/ajmg.c.30013>
31. Boada, R., & Glick, R. (2016). "Klinefelter Syndrome: A Review of the Literature and Clinical Implications." *American Journal of Psychiatry*, 173(5), 498-509. <https://doi.org/10.1176/appi.ajp.2015.15050629>